

Solving some Combinatorial Problems in grid n -ogons

Antonio L. Bajuelos, Santiago Canales, Gregorio Hernández, Ana Mafalda Martins

Abstract — In this paper we study some problems related to grid n -ogons. A grid n -ogon is a n -vertex orthogonal simple polygon, with no collinear edges, that may be placed in a $(n/2) \times (n/2)$ square grid. We will present some problems and results related to a subclass of grid n -ogons, the THIN grid n -ogons, in particular a classification for this subclass of polygons. We follow by presenting the solution of the MINIMUM VERTEX GUARD problem for the Min-Area and for the Spiral grid n -ogons. Finally the solution of the MAXIMUM HIDDEN VERTEX Set problem for THIN grid n -ogons is also presented.

Keywords—Art Gallery Problem, Grid n -ogon, Hidden Set, Orthogonal Polygon

I. INTRODUCTION

IN the field of visibility problems, *guarding* and *hiding* are among the most distinguished and exhaustively studied problems. In visibility problems we are given as input a *simple polygon* (simple closed polygonal curve with its interior). In guarding we need to find a minimum number of guards positioned in the polygon, such that these guards collectively see the whole polygon. Two points in the polygon see each other, if the line segment connecting them lies entirely in the polygon. In hiding, we need to find a maximum number of positions in the polygon, such that no two of these positions see each other.

The guarding problems started during a conference, in 1976, when Victor Klee, posed the following problem, which today is known as the *original art gallery problem*: How many stationary guards are needed to guard an art gallery room with n walls? In the abstract version of this problem, the input is a simple polygon P in the plane, representing the floor plan of the art gallery room and a guard is considered a fixed

point in P with 2π range visibility. A set of guards covers P , if each point of P is seen by at least one guard. Many variations of the original art gallery theorem have been studied over the years, such as: where the guards may be positioned (anywhere or in specific positions, e.g., vertices), what kind of guards are to be used (e.g., stationary guards versus mobile guards) and what assumptions are made for the input polygon (such as being orthogonal) (see [9]). The “opposite” problem of hiding a maximum number of objects from each other in a given simple polygon can have a practical application in computer-games, where a player needs to find and collect or destroy as many objects as possible. Being unable to see the next object while collecting an object makes the game more interesting. Such as the guarding problems, this problem has many variations [2].

In this paper, of the guarding problems, we will consider the MINIMUM VERTEX GUARD (MVG) problem that is the problem of finding the minimum number of guards placed on the vertices (*vertex guards*) needed to guard a given simple polygon. And of the hiding problems, we will consider the MAXIMUM HIDDEN VERTEX SET (MHVS) problem that is the problem of finding the maximum number of vertices of a given simple polygon, such that no two vertices see each other. Both problems are *NP-hard* [3,7]. Important subclasses of polygons are the *orthogonal simple polygons* (simple polygons whose edges meet at right angles). Indeed, they are useful as approximations to polygons; and they arise naturally in domains dominated by Cartesian coordinates, such as raster graphics, VLSI design, or architecture. The MVG and MHVS problems are still *NP-hard* for orthogonal polygons.

This paper has the intention of introducing a particular type of orthogonal polygons - the *grid n -ogons* - that presents sufficiently interesting characteristics that we are studying and formalizing. Of the problems related to grid n -ogons, the visibility problems are the ones that motivate us more, particularly the guarding and hiding problems. The paper is structured as follows: in the next subsection we will introduce some preliminary definitions and useful results. In section 3 we will present some problems and results related to THIN grid n -ogons, in particular a classification for these polygons. In section 4 we will expose some results related to the MVG problem on grid n -ogons and we will study the MHVS problem on THIN grid n -ogons, a subclass of grid n -ogons. Finally, in section 5 we will draw conclusions.

Manuscript received January 24, 2007; Revised received May 12, 2007

This work was supported in part by: the FCT (Portugal) fellowship Grant SFRH/BD/28652/2006, the Grant MEC-HP2005-0137 and CAD-DPI-000235-0505 (Spain) and by CEOC (Portugal) through Program POCTI, FCT, cofinanced by EC fund FEDER and by Acção Integrada N° E-77/06.

Antonio L. Bajuelos, Assistant Professor of Computer Science, Department of Mathematics, University of Aveiro & CEOC, 3810-193 Aveiro, Portugal (phone: (+315)-2343667, fax: (+351)-234382014, e-mail: leslie@ua.pt).

Santiago Canales, Assistant Professor of Computer Science, Escuela Técnica Superior de Ingeniería, Univesidad Potifícia Comillas de Madrid, Spain (e-mail: scanales@upcomillas.es).

Gregorio Hernández, Associate Professor of Computer Science, Facultad de Informática, Universidad Politécnica de Madrid, 28660 Madrid, Spain (e-mail: gregorio@fi.upm.es).

Ana Mafalda Martins, Ph.D student of Computer Science, Department of Mathematics, University of Aveiro & CEOC, 3810-193 Aveiro, Portugal (e-mail: mafalda.martins@ua.pt).

II. CONVENTIONS, DEFINITIONS AND SOME RESULTS

In this paper, the interior and the boundary of a simple polygon P will be denoted by $\text{INT}(P)$ and $\text{BND}(P)$, respectively. And, for convenience, we will assume that the vertices of P are ordered in a counterclockwise (CCW) direction around $\text{INT}(P)$. A vertex of P is called convex if the interior angle between its two incident edges is at most π , otherwise it is called reflex. We use r to represent the number of reflex vertices of P . It has been shown by O'Rourke that $n = 2r + 4$, for every orthogonal simple polygon of n vertices (n -ogon, for short). A *rectilinear cut* of a n -ogon P is a partition of P obtained by extending each edge incident to a reflex vertex of P towards $\text{INT}(P)$ until it hits $\text{BND}(P)$. We denote this partition by $\Pi(P)$ and the number of its pieces by $|\Pi(P)|$. Each piece is a rectangle and so we call it a r -piece. A n -ogon that may be placed in a $(n/2) \times (n/2)$ square grid and that does not have collinear edges is called *grid n -ogon*. We assume that the grid is defined by the horizontal lines $y = 1, \dots, y/2$ and the vertical lines $x = 1, \dots, x/2$ and that its northwest corner is $(1, 1)$. Each grid n -ogon has exactly one edge in every line of the grid. Grid n -ogons that are symmetrically equivalent are grouped [1]. A grid n -ogon Q is called *FAT* iff $|\Pi(Q)| \geq |\Pi(P)|$, for all grid n -ogons P . Similarly, a grid n -ogon Q is called *THIN* iff $|\Pi(Q)| \leq |\Pi(P)|$, for all grid n -ogons P . Let P be a grid n -ogon and r the number of its reflex vertices. In [1] it has been proven that, if P is FAT then $|\Pi(P)| = \frac{3r^2 + 6r + 4}{4}$, for r even and

$|\Pi(P)| = \frac{3(r+1)^2}{4}$, for r odd; if P is THIN then $|\Pi(P)| = 2r + 1$. There is a single FAT grid n -ogon (see Fig. 1(a)); however, THIN grid n -ogons are not unique (see Fig. 1(b)).




Fig. 1: (a) The unique FAT grid n -ogons, for $r = 2, 3$ and 4 ; (b) Two THIN 10-ogons

The area of a grid n -ogon is the number of grid cells in its interior. In [1] it has been proven that for all grid n -ogon P , with $n \geq 8$, $2r + 1 \leq A(P) \leq r^2 + 3$. A grid n -ogon P is a *MAX-AREA* grid n -ogon iff $A(P) = r^2 + 3$ and it is a *MIN-AREA* grid n -ogon iff $A(P) = 2r + 1$. There are *MAX-AREA* grid n -ogons for all n , but they are not unique. However, there is a single *MIN-AREA* grid n -ogon and its form is illustrated in Fig. 2(a). Regarding *MIN-AREA* grid n -ogons, it is obvious that they are *THIN* grid n -ogons, because $|\Pi(P)| = 2r + 1$ holds only for *THIN* grid n -ogons. However, this condition is not sufficient for a *THIN* grid n -ogon to be a *MIN-AREA* grid n -ogon. A grid n -ogon is called a *SPIRAL* grid n -ogon if its

boundary can be divided into a reflex chain and a convex chain. A polygonal chain is called *reflex* if its vertices are all reflex (all except the vertices at the end of the chain) with respect to the interior of the polygon. And, a polygonal chain is called *convex* if its vertices are all convex with respect to the interior of the polygon. In [6] it has been proven that there are *SPIRAL* grid n -ogons, for all $n \geq 6$; however, they are not unique, as we may see this in Fig. 2(b). And it was also proven that every *SPIRAL* grid n -ogon, with $r \geq 1$ reflex vertices, is a *THIN* grid n -ogon.

Given a n -ogon P , we can associate to $\Pi(P)$ a graph, denominated the *dual graph* of $\Pi(P)$ and denoted by $G(P)$, which captures the adjacency relation between pieces of the partition. Each node of the dual graph corresponds to a piece of the partition and its non-oriented edges connect adjacent pieces, i.e., pieces with a common edge. We prove that if P is a *THIN* grid n -ogon then $G(P)$ is a path graph, i.e., a tree with two nodes of vertex of degree 1, called leaves, and the remaining nodes of vertex of degree 2. To prove this result we introduce Lemma 2.1.




Fig. 2: (a) The unique *MIN-AREA* grid n -ogon, for $r = 1, 2$ and 3 ; (b) Two different *SPIRAL* grid 12-ogons (*reflex chain* is bold).

Lemma 2.1: *Let P be a THIN $(n+2)$ -ogon. Then every grid n -ogon that yields P by INFLATE-PASTE (a correct and complete method to generate grid n -ogons, well described in [8]) is also THIN.*

Proposition 2.2: *Let P be a THIN grid n -ogon with r ($r \geq 1$) reflex vertices. Then $G(P)$ is a path graph (see examples in Fig. 3).*

The proof of this proposition is done by induction on r and uses lemma 2.1.




Fig. 3: Three THIN grid 10-ogons and respective dual graphs.

Proposition 2.3: *Let P be a grid n -ogon, with $n > 6$. If P is not THIN then $G(P)$ is not a tree (see example in Fig. 4).*




Fig. 4: A grid 10-ogon and respective dual graph.

Proposition 2.2. establishes that, being P a THIN grid n -ogon, $G(P)$ is a path graph. So, each r -piece of $\Pi(P)$ is adjacent to at most two r -pieces. In this way, each r -piece has at most 2 interior edges. Consequently, we conclude that in $\Pi(P)$ there are 3 types of r -pieces: *Type 1*: with one interior edge and three boundary edges; *Type 2*: with two interior edges not adjacent and two boundary edges not adjacent; *Type 3*: with two adjacent interior edges and two adjacent boundary edges. The r -pieces of the *Type 1* correspond to leaves of $G(P)$ and those of the *Type 2* and *Type 3* correspond to nodes of degree 2. We showed that of the 4 vertices of the r -pieces of *Type 1* three are vertices of P , being two reflex and the other convex, and the other vertex is an interior point of an edge of P . Of the 4 vertices of the r -pieces of *Type 2* two are convex vertices of P and the other two are interior points of edges of P . And finally, of the 4 vertices of the r -pieces of *Type 3* two are vertices of P , being one reflex and the other convex, and the other two are interior points of edges of P (see Fig. 5).




Fig. 5: Types of r -pieces of a THIN grid n -ogon.

Now, we are going to define the *skeleton* of a THIN grid n -ogon. Let P be a THIN grid n -ogon. Since $G(P)$ is a path graph, we can say that P has two “*extremes*”: the r -pieces associated with the leaves of the dual graph. We will denote by *kernel* the extreme that has the horizontal edge with highest y -coordinate. From this graph we can obtain an orthogonal polygonal curve (i.e., a polygonal curve with horizontal or vertical edges) in the following way: we take the centroid of each r -piece, then we connect each one with the centroids of the adjacent r -pieces and, finally, we remove the central vertex of each three aligned vertices, as we can see in Fig. 6. We choose, for the first vertex of this orthogonal curve the kernel's centroid. Therefore it is easy to prove the next result.




Fig. 6: THIN with $r = 4$; on the left is represented its dual graph and on the right its skeleton.

Lemma 2.4: *The skeleton of a THIN grid n -ogon is an orthogonal polygonal curve with $r+2$ vertices.*

III. SOME PROBLEMS RELATED TO THIN GRID N -OGONS

As we have seen in Section 2, on the contrary of the FATS, the THIN n -ogons are not unique. In fact, there are 2 THIN 8-ogon, 30 THIN 10-ogons, 149 THIN 12-ogons, etc. Thus, it is interesting to evidence that the number of THIN grid n -ogons ($|\text{THIN}(n)|$) grows exponentially. Does there exist some expression that relates n to $|\text{THIN}(n)|$? As a step for the resolution of this problem we will first group the THINs into classes. In Section 2 we defined the skeleton of a THIN grid n -ogon. Now, using this concept, we will group THIN grid n -ogons into classes. From the skeleton of the THIN grid n -ogon, we can always represent it by a chain of 0's and 1's, with length r . For that we proceed in the following way: we transverse its skeleton, starting at vertex u_1 , and then we represent each turn left by 1 and each turn right by 0. Now, we will define two operations on these chains: the *complementary operation* and the *inversion operation*.

Definition 3.1: Let c be a chain of 0's and 1's, with length r , i.e., $c = b_1b_2\dots b_r$, where $b_i = 0$ or $b_i = 1$, for $i = 1, 2, \dots, r$. The complementary operation is an operation which takes c as the argument and returns its complementary $c^* = b_1^*b_2^*\dots b_r^*$, where $b_i^* = 1$ if $b_i = 0$ and $b_i^* = 0$ if $b_i = 1$, $i = 1, 2, \dots, r$. The inversion operation is an operation which takes c as the argument and returns its inverse $c^{-1} = b_r b_{r-1} \dots b_1$.

For example, the complementary of the chain $c = 100011$ is the chain $c^* = 011100$ and its inverse is $c^{-1} = 110001$. Easily we can verify that, $(c^*)^{-1} = (c^{-1})^*$, $(c^*)^* = c$ and $(c^{-1})^{-1} = c$.

Proposition 3.2: *Let C_r be the set of all chains, of 0's and 1's, with length r . The relation \sim defined on C_r by $c_1 \sim c_2 \Leftrightarrow c_2 = c_1 \vee c_2 = c_1^{-1} \vee c_2 = c_1^* \vee c_2 = (c_1^*)^{-1}$, is an equivalence relation.*

Consider, now, the quotient set of C_r by \sim , $C_r / \sim = \{[c]_{\sim} : c_1 \in C_r\}$. Note that, each equivalence class has more than one representative. We assume that the representative of each equivalence class always starts by 1.

Proposition 3.3: *Let P_r be the set of all THIN grid n -ogons, with r reflex vertices. The relation \equiv defined on P_r by $P_1 \equiv P_2 \Leftrightarrow c_1 \sim c_2$, where c_1 and c_2 are the chains that represent P_1 and P_2 , respectively, is an equivalence relation.*

The proof of this proposition is trivial. Let $P_r / \equiv = \{[P]_{\equiv} : P_1 \in C_r\}$. Let $P_1, P_2 \in P_r$ and $c_1, c_2 \in C_r$ the chains that represent them, respectively. Note that, P_1 and P_2 belong to the same class (i.e., P_1 and P_2 are *equivalents*) if one of the following conditions is true: (i) $c_1 = c_2$; (ii) $c_2 = c_1^{-1}$; (iii) $c_2 = c_1^*$ or (iv) $c_2 = (c_1^*)^{-1}$. Observe that, geometrically, (ii) can correspond to a horizontal reflection and (iii) to a vertical reflection. In Fig.

7 six THINS with 4 reflex vertices that belong to the same class are illustrated.




Fig. 7: THINS with 4 reflex vertices and respective chains.

We can place the following question: Let c be chain of 0's and 1's with length r , started by 1. Is it always possible to construct a THIN, with r reflex vertices, whose chain that represents it is c ? To answer this question we present the next algorithm:

Algorithm 3.4: Construct a THIN from a chain of 0's and 1's, of length r

Let c be a chain of 0's and 1's, of length r , started by 1.

1. From the chain draw a skeleton ignoring collinearities.
2. Move a vertical sweep line from left to right to eliminate vertical collinearities.
Repeat the previous step until there are no more collinear vertical edges.
3. Move a horizontal sweep line from bottom to top to eliminate horizontal collinearities.
Repeat the previous step until there are no more collinear horizontal edges.

Figure 8 illustrate this algorithm from the chain 1110.




Fig. 8: Constructing the THIN for the chain 1110

Important remarks:

1. In the Step 1, if we construct a skeleton, ignoring the collinearities, we can obtain a skeleton that not correspond to the given chain, for example in the Fig. 9, the chain that represents the constructed Thin is 0010000 (complementary followed by inversion of the given chain).
2. To eliminate collinearities, in step 2 and 3, it is necessary to modify the edge corresponding to the beginning of the polygon. If two edges correspond to the beginning of the polygon, or no edge correspond to

the beginning of the polygon, it is indifferent the one that is modified. Nevertheless, when the polygon has two collinear edges, the choice of the edge is not always indifferent.




Fig. 9: Constructing the THIN from the chain 111011

Anyhow, this algorithm always generates a THIN grid n -ogon whose chain that represents it is equivalent to c . Thus, if the chain that represents the Thin, generated by this algorithm, is c^* ; c^{-1} or $(c^*)^{-1}$, it is enough to make a vertical reflection, a horizontal reflection or a vertical reflection followed by a horizontal reflection, respectively, so that the chain that represents it is exactly c , see Fig.10 for illustration. Thus, this algorithm proves that for each chain of 0's and 1's, with length r , started by 1, there is a Thin grid n -ogon with r reflex vertices represented by it.




Fig. 10: (a) The chain that represents the THIN after the reflections is $c = 111011$; (b) The chain that represents the THIN after the reflections is $c = 1001$.

Based on the previous reasoning and definitions it is not difficult to prove the following result.

Proposition 3.5: The correspondence $f: P_r/\cong \rightarrow C_r/\sim$ defined by $f([P_1]) = [c_1]$, where $c_1 \in C_r$ is the chain that represents $P_1 \in P_r$, which is a representative of the class $[P_1]$, is a bijective function.

Now, we may show the next result that allows us to count the number of classes of THIN grid n -ogons with r reflex vertices.

Proposition 3.6: *The number of classes of THIN grid n -ogons with r reflex vertices ($r \geq 2$) is equal to:*

$$\begin{cases} 2^{r-2} + 2^{\frac{1}{2}(r-3)}, & \text{if } r \text{ is odd} \\ 2^{r-2} + 2^{\frac{1}{2}(r-2)}, & \text{if } r \text{ is even} \end{cases}$$

Proof. By proposition 3.5 we can conclude that $|P_{r/\equiv}| = |C_{r/\sim}|$, so we just have to calculate $|C_{r/\sim}|$. The cardinal of C_r is 2^r and the number of symmetrical chains ($c = c^{-1}$), with length r , is $2^{\lceil r/2 \rceil}$. If a chain c is symmetrical, then its equivalence class is constituted by two chains, c and c^* . If a chain c is not symmetrical, to find the cardinal of its class, we have to distinguish two cases: r odd and r even. If r is odd, all the chains have 4 equivalent chains: c , c^{-1} , c^* and $(c^*)^{-1}$ (for example: $c = 11010, 01011, 00101$ and 10100). If r is even, there are chains that have 4 equivalent chains (e.g., $c = 1110$) and chains that only have 2 equivalent chains; this case happens when $c^* = c^{-1}$ (e.g., for the chain $c = 1100$, $c^* = c^{-1} = 0011$). Let us now count the number of equivalence classes. If r is odd, the number of equivalence classes of the symmetrical chains (SC) is $\frac{1}{2}|SC| = \frac{1}{2} \left(2^{\frac{r+1}{2}} \right) = 2^{\frac{r-1}{2}}$ and the number of

equivalence classes of the non symmetrical chains (NSC) is $\frac{1}{4}|NSC| = \frac{1}{4} \left(2^r - 2^{\frac{r+1}{2}} \right) = 2^{r-2} - 2^{\frac{r-3}{2}}$. Thus, if r is odd, the total

number of equivalence classes is $2^{r-2} + 2^{\frac{r-1}{2}} - 2^{\frac{r-3}{2}} = 2^{r-2} + 2^{\frac{1}{2}(r-3)}$. If r is even, the number of equivalence classes of symmetrical chains is equal to $\frac{1}{2}|SC| = \frac{1}{2} \left(2^{\frac{r}{2}} \right) = 2^{\frac{1}{2}(r-2)}$. The

number of equivalence classes of non symmetrical chains constituted by two chains (for example, the classes of the chains $101010, 1100, 110100, \dots$) is $\frac{1}{2} \left(2^{\frac{r}{2}} \right) = 2^{\frac{1}{2}(r-2)}$. In fact,

to obtain $c^* = c^{-1}$, the second half of the chain is completely determined by the first half. Therefore, the cardinal of these classes is half of the number of chains of this type. And, the number of equivalence classes of non symmetrical chains constituted by four chains is $\frac{1}{4}|All - Symmetric - (Chains with c^* = c^{-1})| = \frac{1}{4} \left(2^r - 2^{\frac{r}{2}} - 2^{\frac{r}{2}} \right) = 2^{r-2} - 2^{\frac{1}{2}(r-2)}$. Thus, if r is even, in the total, the number of equivalence classes is equal to $2^{r-2} + 2^{\frac{1}{2}(r-2)}$ \square

However, there are still some open problems to solve, such as: How many elements THIN grid n -ogons does each class

have? Will it be possible to find an algorithm that generates all THIN grid n -ogons of the same class?

Note that, solving the first problem we also solve the initial problem, that is: Does there exist some expression that relates n to $|THIN(n)|$?

IV. SOME VISIBILITY PROBLEMS ON GRID n -OGONS

Of the problems related to grid n -ogons, the guarding and hiding problems are the ones that motivate us more, particularly the MVG and MHVS problems. Since THIN and FAT n -ogons are the classes for which the number of r -pieces is minimum and maximum, we think that they can be representative of extremal behavior. Besides that they are used experimentally to evaluate approximate methods of resolution of the MVG problem, so we started with them. We have already proven that to guard any FAT grid n -ogon it is always sufficient two $\pi/2$ vertex guards (vertex guards with $\pi/2$ range visibility) and established where they must be placed [4]. However, THIN grid n -ogons are much more difficult to guard, in spite of having much fewer r -pieces than FATS. Besides, they are not unique, so we tried to characterize structural properties of classes of THINS that allow for simplifying the problem's study. Up to now the only quite characterized subclasses are the MIN-AREA and the SPIRAL grid n -ogons. We proved that to guard any MIN-AREA and SPIRAL grid n -ogon $\lceil n/6 \rceil$ and $\lfloor n/4 \rfloor$ vertex guards are necessary, respectively. Moreover, we showed where those guards could be placed [5, 6].

MAXIMUM HIDDEN VERTEX Set Problem on THIN grid n -ogons

Given a simple polygon, P , and a subset of vertices of P , HV , we say that HV is a *hidden vertex set* if no two vertices in HV see each other. The MAXIMUM HIDDEN VERTEX SET problem on a simple polygon asks for an hidden vertex set, HV , of maximum cardinality. We will call the elements of HV *hidden vertices*. Shermer [7] proved that the size of the MAXIMUM HIDDEN VERTEX SET of a n -ogon is at most $(n-2)/2$. This tight bound is achieved in staircase polygons. We will show that, given a THIN grid n -ogon the maximum cardinality of a hidden vertex set is $\lceil n/4 \rceil$.

Let P be a THIN grid n -ogon and $S = u_1 u_2 \dots u_{n/2}$ its skeleton. Let us assume, without loss of generality that the first edge of S , $[u_1 u_2]$, is horizontal and that u_2 is to the right of u_1 . Note that, the boundary of P consists of two joined polygonal chains, c_1 and c_2 , "parallel" to S , where the first edge of c_1 is a bottom edge and the first edge of c_2 is a top edge. Note that, c_1 and c_2 can be expressed as ordered sequences of vertices $c_1 = v_1^1 v_2^1 \dots v_{n/2}^1$ and $c_2 = v_1^2 v_2^2 \dots v_{n/2}^2$, where v_i^1 denotes the i^{th} vertex of c_1 and v_i^2 denotes the i^{th} vertex of c_2 (see Fig. 11). This way, $BND(P) = c_1 \cup [v_{n/2}^1 v_{n/2}^2] \cup c_2 \cup [v_1^2 v_1^1]$. Observe, also, that, if we transverse S , starting at vertex u_1 , c_1 is always on the right of S and c_2 on the left.




Fig. 11: Two THINS grid n -ogons, its skeletons and the chains c_1 and c_2 (c_1 in bold).

To each vertex of the skeleton we correspond two vertices of the polygon, one in c_1 and another one in c_2 . That is, to $u_i \in S$, we correspond the vertices $v_i^1 \in c_1$ and $v_i^2 \in c_2$. And to each edge of the skeleton we correspond two parallel edges of the polygon, one in c_1 and another one in c_2 . That is, to $[u_i, u_{i+1}] \in S$, we correspond the edges $[v_i^1, v_{i+1}^1] \in c_1$ and $[v_i^2, v_{i+1}^2] \in c_2$. Note that, by construction of the skeleton, we can easily see that any point of $[v_i^1, v_{i+1}^1]$ sees any point of $[v_i^2, v_{i+1}^2]$.

Now, for each $u_{2k-1} \in S$ with $k = 1, \dots, \lceil n/4 \rceil$, we mark a hidden vertex in P , in the following way: for $k = 1$ we mark v_1^1 ; for $k \neq 1$ we mark v_{2k-1}^1 or v_{2k-1}^2 , depending if v_{2k-2}^1 is reflex or convex, respectively (see Fig. 12, for illustration).




Fig. 12: Two THINS n -ogons and marked hidden vertices (c_1 in bold).

Note that, the $\lceil n/4 \rceil$ marked vertices form a hidden vertex set, since each time that we mark a new vertex as hidden we can guarantee that it does not see none of the vertices that previously had been marked as hidden. In fact, for $k = 1$ it is trivial. For $k \neq 1$, we have two cases, depending if v_{2k-2}^1 is reflex (*Case 1*) or convex (*Case 2*), as we can see in Fig.13.




Fig. 13: In *Case 1* the vertex v_{2k-2}^1 is reflex and in *Case 2* the vertex v_{2k-2}^1 is convex.

In *Case 1* the vertex that is marked as hidden is the vertex v_{2k-1}^1 and in *Case 2* is the vertex v_{2k-1}^2 . In both cases the marked vertex does not see none of the already marked as hidden, since of the already “visited” vertices this one only sees v_{2k-1}^1 and v_{2k-1}^2 (see Fig.14).




Fig. 14: The shaded zones are not visible by the marked vertices.

Observe that if we mark as hidden the vertices v_{2k-1}^2 (in *Case 1*) and v_{2k-1}^1 (in *Case 2*) we can't guarantee that they don't see any of the vertices already marked as hidden, since they see more backwards (see Fig. 15).




Fig. 15: The shaded zones are not visible by the marked vertices.

Therefore it is easy to prove the next result.

Lemma 4.1: For any THIN grid n -ogon there is an hidden vertex HV and $|HV| = \lceil n/4 \rceil$.

And then using this Lemma we prove our main result.

Theorem 4.2: Let P be a THIN grid n -ogon. The maximum cardinality of an hidden vertex set in P is $\lceil n/4 \rceil$.

V. CONCLUSION

We presented some results related to grid n -ogons. Of the hiding problems related to the grid n -ogons, it is the MHVS problem that motivates us more. We proved that the maximum cardinality of an hidden vertex set in a THIN n -ogon is $\lceil n/4 \rceil$. Moreover, we established a possible positioning for those hidden vertices. We also established a possible classification for THIN n -ogon, as a step to launch an expression that relates n to $\lceil \text{THIN}(n) \rceil$.

ACKNOWLEDGMENT

We wish to thank Inês Pereira Matos from the University of Aveiro for helpful discussions about the problem of classification of THIN grid n -ogons.

REFERENCES

- [1] A. L. Bajuelos, A. P. Tomás: Partitioning Orthogonal Polygons by Extension of All Edges Incident to Reflex Vertices: lower and upper bounds on the number of pieces. Proc. of ICCSA 2004, LNCS 3045, Springer-Verlag (2004) 127-136.
- [2] S. Eidenbenz: In-approximability of Finding Maximum Hidden Sets on Polygons and Terrains. Computational Geometry 21 (2002) 139–53.
- [3] D. Lee, A. Lin: Computational Complexity of Art Gallery Problems. IEEE Transactions on Information Theory IT-32 (1986) 276-282.
- [4] A. M. Martins, A. Bajuelos: Some Properties of Fat and Thin grid n -ogons. in Wiley-VCH Verlag (Eds): Proc. of ICNAAM 2005 (2005) 361-365.
- [5] A. M. Martins, A. Bajuelos: Characterizing and Covering some Classes of Orthogonal polygons. In Proc. ICCS 2006, LNCS 3992, Springer-Verlag (2006) 255-262.
- [6] A. M. Martins, A. Bajuelos: Vertex Guards in a Subclass of Orthogonal Polygons. International Journal of Computer Science and Network Security (IJCSNS), Vol. 6 No. 9, September 2006.
- [7] T. Shermer: Hiding people in polygons, Computing 42 (1989) 109–131.
- [8] A. P. Tomás, A. Bajuelos: Quadratic-Time Linear-Space Algorithms for Generating Orthogonal Polygons with a Given Number of Vertices. In Proc. of ICCSA 2004, LNCS 3045, Springer-Verlag (2004) 117-126.
- [9] J. Urrutia: Art Gallery and Illumination Problems. In J.-R. Sack and J. Urrutia (editors), Handbook of Computational Geometry, Elsevier (2000).