

- [36] C. W. J. Beenakker, "Colloquium: Andreev reflection and Klein tunneling in graphene," *Rev. Mod. Phys.*, vol. 80, pp. 1337–1354, 2008, DOI: 10.1103/RevModPhys.80.1337.
- [37] A. F. Young, P. Kim, "Quantum interference and Klein tunnelling in graphene heterojunctions," *Nature Phys.*, vol. 5, pp. 222–226, 2009, DOI: 10.1038/nphys1198.
- [38] V. Nam Do, V. Hung Nguyen, P. Dollfus, and A. Bournel, "Electronic transport and spin-polarization effects of relativistic-like particles in mesoscopic graphene structures," *J. Appl. Phys.*, vol. 104, p. 063708, 2008, DOI: 10.1063/1.2980045.
- [39] V. Hung Nguyen, A. Bournel, and P. Dollfus, "Resonant tunneling structures based on epitaxial graphene on SiC," *Semicond. Sci. Technol.*, vol. 26, p. 125012, 2011, DOI: 10.1088/0268-1242/26/12/125012.
- [40] P. E. Allain and J. N. Fuchs, "Klein tunneling in graphene: Optics with massless electrons," *Eur. Phys. J. B*, vol. 83, pp. 301–317, 2011, DOI: 10.1140/epjb/e2011-20351-3.
- [41] I. Rodríguez-Vargas, J. Madrigal-Melchor, and O. Oubram, "Resonant tunneling through double barrier graphene systems: A comparative study of Klein and non-Klein tunneling structures," *J. Appl. Phys.*, vol. 112, p. 073711, 2012, DOI: 10.1063/1.4757591.
- [42] J. M. Pereira Jr., F. M. Peeters, A. Chaves, and G. A. Farias, "Klein tunneling in single and multiple barriers in graphene," *Semicond. Sci. Technol.*, vol. 25, p. 033002, 2010, DOI: 10.1088/0268-1242/25/3/033002.
- [43] C.-H. Park, Y.-W. Son., L. Yang, M. L. Cohen, and S. G. Louie, "Electron Beam Supercollimation in Graphene Superlattices," *Nano Lett.*, vol. 8, pp. 2920–2924, 2008, DOI: 10.1021/nl801752r.
- [44] D. Dragoman and M. Dragoman, "Negative differential resistance of electrons in graphene barrier," *Appl. Phys. Lett.*, vol. 90, p. 143111, 2007, DOI: 10.1063/1.2719670.
- [45] R. Zhu and Y. Guo, "Shot noise in the graphene-based double-barrier structures," *Appl. Phys. Lett.*, vol. 91, p. 252113, 2007, DOI: 10.1063/1.2825571.
- [46] T. Tudorovskiy, K. J. A. Reijnders, and M. I. Katsnelson, "Chiral tunneling in single-layer and bilayer graphene," *Phys. Scr.*, vol. T146, p. 014010, 2012, DOI: 10.1088/0031-8949/2012/T146/014010.
- [47] N. T. T. Nguyen, D. Q. To, and V. L. Nguyen, "A model for ballistic transport across locally gated graphene bipolar junctions," *J. Phys. Condens. Matter*, vol. 26, p. 015301, 2014, DOI: 10.1088/0953-8984/26/1/015301.
- [48] H. C. Nguyen and V. L. Nguyen, "Tunneling of Dirac electrons through one-dimensional potentials in graphene: a T -matrix approach," *J. Phys.: Condens. Matter*, vol. 21, p. 045305, 2009, DOI: 10.1088/0953-8984/21/4/045305.
- [49] L. Brey and H. A. Fertig, "Emerging zero modes for graphene in a periodic potential," *Phys. Rev. Lett.*, vol. 103, p. 046809, 2009, DOI: 10.1103/PhysRevLett.103.046809.
- [50] P. Burset, A. Levy Yeyati, L. Brey, and H. A. Fertig, "Transport in superlattices on single-layer graphene," *Phys. Rev. B*, vol. 83, p. 195434, 2011, DOI: 10.1103/PhysRevB.83.195434.
- [51] M. Barbier, P. Vasilopoulos, and F. M. Peeters, "Extra Dirac points in the energy spectrum for superlattices on single-layer graphene," *Phys. Rev. B*, vol. 81, p. 075438, 2010, DOI: 10.1103/PhysRevB.81.075438.
- [52] M. Barbier, P. Vasilopoulos, and F. M. Peeters, "Single-layer and bilayer graphene superlattices: collimation, additional Dirac points and Dirac lines," *Phil. Trans. R. Soc. A*, vol. 368, pp. 5499–5524, 2010, DOI: 10.1098/rsta.2010.0218.
- [53] S. Dubey, V. Singh, A. K. Bhat, P. Parikh, S. Grover, R. Sensarma, V. Tripathi, K. Sengupta, and M. M. Deshmukh, "Tunable superlattice in graphene to control the number of Dirac points," *Nano Lett.*, vol. 13, pp. 3990–3995, 2013, DOI: 10.1021/nl4006029.
- [54] G. J. Xu, X. G. Xu, B. H. Wu, J. C. Cao, and C. Zhang, "The resonant tunneling through a graphene multiquantum well system," *J. Appl. Phys.*, vol. 107, p. 123718, 2010, DOI: 10.1063/1.3445782.
- [55] Y. O. Klymenko and O. Shevtsov, "Quantum transport in armchair graphene ribbons: analytical tight-binding solutions for propagation through step-like and barrier-like potentials," *Eur. Phys. J. B*, vol. 69, pp. 383–388, 2009, DOI: 10.1140/epjb/e2009-00178-3.
- [56] Y. O. Klymenko, L. Malysheva, and A. Onipko, "Electron transmission through step- and barrier-like potentials in graphene ribbons," *Phys. Stat. Sol. (b)*, vol. 245, pp. 2181–2184, 2008, DOI: 10.1002/pssb.200879637.
- [57] A. Cresti, G. Grosso, and G. Pastori Parravicini, "Numerical study of electronic transport in gated graphene ribbons," *Phys. Rev. B*, vol. 76, p. 205433, 2007, DOI: 10.1103/PhysRevB.76.205433.
- [58] O. Roslyak, A. Iurov, G. Gumbs, and D. Huang, "Unimpeded tunneling in graphene nanoribbons," *J. Phys.: Cond. Matt.*, vol. 22, p. 165301, 2010, DOI: 10.1088/0953-8984/22/16/165301.
- [59] J. H. Oh and D. Ahn, "Klein Tunneling Through an Oblique Barrier in Graphene Ribbons," *J. Korean Phy. Soc.*, vol. 59, pp. 2786–2791, 2011, DOI: 10.3938/jkps.59.2786
- [60] E. D. Herbschleb, R. K. Puddy, P. Marconcini, J. P. Griffiths, G. A. C. Jones, M. Macucci, C. G. Smith, and M. R. Connolly, "Direct imaging of coherent quantum transport in graphene p-n-p junctions," *Phys. Rev. B*, vol. 92, p. 125414, 2015, DOI: 10.1103/PhysRevB.92.125414.
- [61] P. Marconcini and M. Macucci, "Numerical analysis of the resistance behavior of an electrostatically-induced graphene double junction," *J. Comput. Electron.*, vol. 14, p. 653–660, 2015, DOI: 10.1007/s10825-015-0701-5.
- [62] P. Marconcini and M. Macucci, "Approximate calculation of the potential profile in a graphene-based device," *IET Circ. Device Syst.*, vol. 9, pp. 30–38, 2015, DOI: 10.1049/iet-cds.2014.0003.
- [63] D. Logoteta, P. Marconcini, C. Bonati, M. Fagotti, and M. Macucci, "High-performance solution of the transport problem in a graphene armchair structure with a generic potential," *Phys. Rev. E*, vol. 89, p. 063309, 2014, DOI: 10.1103/PhysRevE.89.063309.
- [64] H. B. Nielsen and M. Ninomiya, "A no-go theorem for regularizing chiral fermions," *Phys. Lett. B*, vol. 105, pp. 219–223, 1981, DOI: 10.1016/0370-2693(81)91026-1.
- [65] P. Marconcini, "The role of the choice of the physical model in the optimization of nanoelectronic device simulators," *International Journal of Circuits, Systems and Signal Processing*, vol. 7, pp. 173–180, 2013.
- [66] M. Fagotti, C. Bonati, D. Logoteta, P. Marconcini, and M. Macucci, "Armchair graphene nanoribbons: \mathcal{PT} -symmetry breaking and exceptional points without dissipation," *Phys. Rev. B*, vol. 83, p. 241406(R), 2011, DOI: 10.1103/PhysRevB.83.241406.
- [67] P. Marconcini, D. Logoteta, and M. Macucci, "Sinc-based method for an efficient solution in the direct space of quantum wave equations with periodic boundary conditions," *J. Appl. Phys.*, vol. 114, p. 173707, 2013, DOI: 10.1063/1.4828875.
- [68] S. Datta, *Electronic Transport in Mesoscopic Systems*, Cambridge, UK: Cambridge University Press, 1997, ISBN: 9780521599436.
- [69] R. Landauer, "Spatial variation of currents and fields due to localized scatterers in metallic conduction," *IBM J. Res. Dev.*, vol. 1, pp. 223–231, 1957, DOI: 10.1147/rd.13.0223
- [70] M. Büttiker, Y. Imry, R. Landauer, and S. Pinhas, "Generalized many-channel conductance formula with application to small rings," *Phys. Rev. B*, vol. 31, pp. 6207–6215, 1985, DOI: 10.1103/PhysRevB.31.6207.
- [71] M. Büttiker, "Scattering theory of thermal and excess noise in open conductors," *Phys. Rev. Lett.*, vol. 65, pp. 2901–2904, 1990, DOI: 10.1103/PhysRevLett.65.2901.
- [72] E. Anderson, Z. Bai, C. Bischof, S. Blackford, J. Demmel, J. Dongarra, J. Du Croz, A. Greenbaum, S. Hammarling, A. McKenney, and D. Sorensen, *LAPACK Users' Guide*. Philadelphia, PA: Society for Industrial and Applied Mathematics, 1999, ISBN: 0898714478.
- [73] V.-Hung Nguyen, "Electronic transport and spin-polarization effects in graphene nanostructures," Ph.D. Thesis, Université de Paris-Sud, Ecole doctorale Sciences et Technologies de l'Information, des Télécommunications et des Systèmes, Orsay, Essonne, France, 2010.



Paolo Marconcini Paolo Marconcini received the Master's (summa cum laude) degree in Electrical Engineering and the Ph.D. degree in Electrical and Computer Engineering from the Università di Pisa, Pisa, Italy, in 2002 and 2006, respectively. From 2006, he has been a Post-Doctoral Researcher and then Tenure-Track Researcher with the Università di Pisa, where he is now Associate Professor. He has authored over 100 papers in international journals and proceedings and has been involved in over ten national and international research projects. His

current research interests include the study of nanoelectronic devices based on semiconductor heterostructures, nanowires, carbon nanotubes, graphene and two-dimensional materials, and of electronic systems for transportation.

Creative Commons Attribution License 4.0 (Attribution 4.0 International, CC BY 4.0)

This article is published under the terms of the Creative Commons Attribution License 4.0

https://creativecommons.org/licenses/by/4.0/deed.en_US