salinity buildup and transpiration. *Acta Horiculturae* 573 :97-106.

[5] Beltrão, J.; & Ben Asher, J. 1997. The effect of salinity on corn yield using CERES-Maize model. *Irrigation and Drainage Systems* 11:15-28.

[6] Hamdy, A. 2002. Sustainable use and management of non-conventional water resources in the arid regions. *Acta Horiculturae* 573:159-174.

[7] Cuartero; J.; Pulido, J. M.; Gómez-Guillaumón, M.L.; & Alvarez, M. 2002. Salt removal potential of barley, alfalfa, *Atriplex patula* and *A. prostate. Acta Horiculturae* 573:387-396.

[8] Beltrão, J.; Brito, J.; Neves, M.A.; & Seita, J. 2009. Salt Removal Potential of Turfgrasses in Golf Courses in the Mediterranean Basin. *WSEAS Transactions on Environment and Development* 5 (5):394-403.

[9] Hamidov, A.; Khaydarova, Khamidov, M.; Neves, M.A.;.& Beltrao, J. 2007. *Apocynum Lancifolium* and *Chenopodium Album*.- potencial species to remediate saline soils. *WSEAS Transactions on Environment and Development* 7(3):123-128.

[10] Glenn, E.P; Brown, J.J.; & Blumwald, E. 1999. Salt tolerance and crop potential of halophytes. *Critical Reviews in Plant Sciences* 18:227-255.

[11] Nasr, Z.; & Ben Mechlia, N. 2002. Deficit irrigation to reduce salinization in an apple orchard. *Acta Horticulturae* 573:283-287.

[12] Beltrão, J.; Gamito, P.; Guerrero, Arsénio, A. & Brito, J.M.C.1999. Grass response to municipal wastewater as compared to nitrogen and water application. In: D. Anaç & P. Martin-Prével (Eds.). Improved Crop quality by Nutrient management, Chapter 60: 263-266. Kluwer Academic Publishers.

[13] Shaaban, M.M.; & El-Fouly, M.M. 2002. Nutriente contents and salt removal potential of some wiold plants grown in salt affected soils. *Acta Horiculturae* 573:377-385.

[14] Cuartero, J.; Pulido, J.M. and Gomez-Guilamon La Mayora, M.L. 2002. Salt Removal Potential of Barley, Alfalfa, Atriplex patula and A. prostrata. *Acta Horiculturae* 573:387-391.

[15] Anac, S; Kukul, Y. and Anac, D. 2005. Salt removing crops as a phytoremediation technique. Internetional meeting on phytodepuration.

[16] Kiliç, C.C., Kukul, Y., Anac, D., 2008. Performance of purslane (Portulaca oleracea L.) as a salt-removing crop. Agricultural water management 95 (7) 854-858.

[17] Neves, M.A.; Miguel, M.G.; Marques, C.; Panagopoulos, T.; Beltrão, J. 2006. Response of Tetragonia tetragonoides (Pallas) Kuntze to the combined effects of salts and nitrogen. *WSEAS Transactions on Environment and Development* 2(4):470-474.

[18] Beltrão, J.; Jesus, S.B.; Trindade, D.; Miguel, M.G.; Neves, M.A.; Panagopoulus, T. & Ben Asher, J. 2002. Combined effects of salts and nitrogen on the yield function of Lettuce. *Acta Horticulturae* 573:363-368.

[19] Neves, M.A.; Miguel, M.G.; Marques, C.; Panagopoulos, T.; Beltrão, J. 2008. The combined effects of salts and calcium on growth and mineral accumulation on *Tetragonia tetragonioides*. *WSEAS Transactions on Environment and Development* 4(1):1-5

[20] Hamidov, A.; Beltrao, J.; Costa, C.; Khaydarova, V.; Sharipov, S. 2007. Environmentally Useful Technique – *Portulaca oleracea* Golden Purslane as a Salt Removal Species. *WSEAS Transactions on Environment and Development* 2 (4):117-122.

[21] ADB supported project RETA. 2008. Enabling communities in the Aral Sea Basin to Combat land and water resource degradation through the creation of bright spots supported Report nº 6208:54-64.

[22] Piper, C.S. Soil and Plant analyses. Inter- Science Publishers Inc., New York, 1945.

[23] Chapman, H.D., Pratt, P.F. Methods of analyses for soils, plants and waters. University of California, Division of Agriculture Science, Riverside. 1961.

[24] SPSS 11.0. 2002. SPSS Inc., Chicago, Illinois, U.S.A.