

# Spectrum of Fibonacci and Lucas numbers

Asker Ali Abiyev

**Abstract** It has been achieved polynomial function, depending on arguments  $a + b$  and  $ab$  arguments of expression  $a^n + b^n$  for the biggest and smallest numbers which are in the centre of natural geometrical figures (line, square, cube, ..., hypercube). The coefficients of this polynomial are defined from triangle tables, written by special algorithm by us. The sums of the numbers in each row of the triangles make Lucas and Fibonacci sequences. New formulae for terms of these sequences have been suggested by us (Abiyev's theorem). As the coefficients of the suggested polynomial are spectrum of Fibonacci and Lucas numbers they will opportunity these number's application field to be enlarged.

**Keywords:** Lucas, Fibonacci, sequences, identity, Binet formula, polynomial.

## 1. INTRODUCTION

The binomial expression  $a^n \pm b^n$  is one of the most used expressions in algebra. It has evidently been seen up to now that one of the powers of  $a$  and  $b$  coefficients increases and another one decreases in the identity of this expression [1]. The following formula can serve as an example.

$$a^n - b^n = (a - b)(a^{n-1} + a^{n-2}b + a^{n-3}b^2 + \dots + ab^{n-2} + b^{n-1}) \quad (1)$$

When  $a$  and  $b$  letters are conjugate expressions, algebraic calculus become more difficult by using (1) identity. As the sum and multiplication of the conjugate expressions are real numbers, writing the right side of (1) identity as a function depending on  $a + b$  and  $ab$  arguments simplifies the calculus. This function can be easily written for small values of  $n$ :

$$a^3 + b^3 = (a + b) \left[ (a + b)^2 - 3ab \right]$$

$$a^3 - b^3 = (a - b) \left[ (a + b)^2 - ab \right]$$

But there are no formulae for big values of  $n$  in literature. The purpose of the article is to write new identities for the binomial expression  $a^n \pm b^n$  and study the fields of their application.

## 2. $F[(a + b), ab]$ EXPRESSION

If set of natural numbers  $\{1, 2, 3, \dots, n\}$ ,  $\{1, 2, 3, \dots, n^2\}$ ,  $\{1, 2, 3, \dots, n^3\}$ ,  $\{1, 2, 3, \dots, n^4\}$ , ...,  $\{1, 2, 3, \dots, n^p\}$  is written accordingly in line, square, cube, ..., hypercube of  $p$  order one after another, the amount of numbers in their centres is defined with  $q=2^p$  formula. Let's mark the maximal number with  $A_p$  and minimal number with  $B_p$ . Where,  $n$ - even number and order of geometrical figure,  $p$ - dimension of space. If  $p=3$  and  $n=6$ , it is cube of 6 order in space of 3 dimension.

The following formulae  $A_p(n)$  and  $B_p(n)$  achieved after doing some algebraic operations by using numbers of figures' centres for  $p=1$ ,  $p=2$ ,  $p=3$ ,  $n=6$  and  $n=8$ :

$$A_p(n) = \frac{n}{2} \left[ n^{p-1} + (n^{p-2} + n^{p-3} + \dots + 1) \right] + 1 \quad (2)$$

$$B_p(n) = \frac{n}{2} \left[ n^{p-1} - (n^{p-2} + n^{p-3} + \dots + 1) \right] \quad (3)$$

Let's calculate  $A_2(6)$ ,  $B_2(6)$ ,  $A_3(6)$ ,  $B_3(6)$  by these formulae:

$$A_2(6) = \frac{6}{2} [6 + 1] + 1 = 22 \quad ; \quad B_2(6) = \frac{6}{2} [6 - 1] = 15 \quad ;$$

$$A_3(6) = \frac{6}{2} [6^2 + (6 + 1)] + 1 = 3 \cdot 43 + 1 = 130 \quad ,$$

$$B_3(6) = \frac{6}{2} [6^2 - (6 + 1)] = 3 \cdot 29 = 87 \quad .$$

And now let's find out sum and product of  $A_p(n)$  and  $B_p(n)$ :

$$A_p(n) + B_p(n) = \frac{n}{2} \left[ n^{p-1} + (n^{p-2} + n^{p-3} + \dots + 1) \right] + 1 + \frac{n}{2} \left[ n^{p-1} - (n^{p-2} + n^{p-3} + \dots + 1) \right] = n^p + 1 \quad (4)$$

$$A_p(n) \cdot B_p(n) = \frac{n}{2} \left\{ n^{p-1} \left( \frac{n^p}{2} + 1 \right) - (n^{p-2} + n^{p-3} + \dots + 1) \left[ \frac{n}{2} (n^{p-2} + n^{p-3} + \dots + 1) + 1 \right] \right\} \quad (5)$$

By these formula let's calculate  $A_2(6) + B_2(6)$ ,  $A_2(6) \cdot B_2(6)$  and  $A_3(6) + B_3(6)$ ,  $A_3(6) \cdot B_3(6)$ :

$$A_3(6) + B_3(6) = 6^3 + 1 = 217 = 130 + 87; \quad A_3(6) \cdot B_3(6) = 3 \{ 36 \cdot 109 - 7 \cdot [3 \cdot 7 + 1] \} = 3 \cdot 3770 = 11 \cdot 310 = 130 \cdot 87$$

(4) and (5) formulae have been proved by mathematical induction method.

After using Binomial theorem and doing some algebraic operations, we get the following equalities for

$A_p^{p+1} + B_p^{p+1} = F[(A_p + B_p), A_p B_p]$  expression: if p- odd

$$A_p^{p+1} + B_p^{p+1} = E_{p+1}^1 (A_p + B_p)^{p+1} - A_p B_p \times \left[ E_{p+1}^2 (A_p + B_p)^{p-1} - A_p B_p \left[ E_{p+1}^3 (A_p + B_p)^{p-3} - \dots - A_p B_p \left( E_{p+1}^{\frac{p+1}{2}} (A_p + B_p)^2 - E_{p+1}^{\frac{p+3}{2}} A_p B_p \right) \dots \right] \right] \quad (6)$$

if p- even

$$\frac{A_p^{p+1} + B_p^{p+1}}{A_p + B_p} = T_{p+1}^1 (A_p + B_p)^p - A_p B_p \times \left[ T_{p+1}^2 (A_p + B_p)^{p-2} - A_p B_p \left[ T_{p+1}^3 (A_p + B_p)^{p-4} - \dots - A_p B_p \left( T_{p+1}^{\frac{p}{2}} (A_p + B_p)^2 - T_{p+1}^{\frac{p+1}{2}} A_p B_p \right) \dots \right] \right] \quad (7)$$

$E_{p+1}^1, E_{p+1}^2, \dots, E_{p+1}^{\frac{p+3}{2}}$  and  $T_{p+1}^1, T_{p+1}^2, \dots, T_{p+1}^{\frac{p+1}{2}}$  coefficients in (6) and (7) expressions are found from the Table 1.

Table1, drawn with special algorithm by us [2]. Where  $E_{p+1}^1 = T_{p+1}^1 = 1$

					E, T	p+1
					1	1
				1	2	2
			1	1	3	3
		1	1	4	2	4
		1	1	5	5	5
	1	6	1	9	2	6
	1	7	1	14	7	7
1	8	20	1	16	2	8
1	9	27	1	30	9	9
1	10	35	50	25	2	10
...	...	...	...	...	...	...

If to use (2) and (4) expressions, a new equality is found for  $A_p B_p$  product:

$$A_p B_p = \frac{n^p}{2} \left( \frac{n^p}{2} + 1 \right) - \frac{n}{2} (n^{p-2} + n^{p-3} + \dots + 1) \left[ \frac{n}{2} (n^{p-2} + n^{p-3} + \dots + 1) + 1 \right] = \frac{(s_p - 1)(s_p + 1)}{4} - (A_{p-1} - 1) A_{p-1} = D_p \quad (8)$$

If to use (4) and (8) equalities, (6) and (7) will be as following:

$$A_p^{p+1} + B_p^{p+1} = E_{p+1}^1 S_p^{p+1} - D_p \left( E_{p+1}^2 S_p^{p-1} - D_p \left( E_{p+1}^3 S_p^{p-3} - \dots - D_p \left( E_{p+1}^{\frac{p+1}{2}} S_p^2 - E_{p+1}^{\frac{p+3}{2}} D_p \right) \dots \right) \right) \quad (9)$$

$$A_p^{p+1} + B_p^{p+1} = s_p \left( T_{p+1}^1 s_p^p - D_p \left( T_{p+1}^2 s_p^{p-2} - D_p (T_{p+1}^3 s_p^{p-4} - \dots - D_p (T_{p+1}^{\frac{p}{2}} s_p^2 - T_{p+1}^{\frac{p}{2}+1} D_p) \dots) \right) \right) \quad (10)$$

$$A_1^2 + B_1^2 = E_2^1 s_1^2 - E_2^2 D_1 = s_1^2 - 2D_1 = (n+1)^2 - 2 \frac{(n+2)n}{4} - (A_0 - 1)A_0 = \frac{(n+1)^2 + 1}{2} = \left(\frac{n}{2} + 1\right)^2 + \left(\frac{n}{2}\right)^2 \quad (11)$$

Let's calculate (10) and (11) equalities for p=1, p=2, p=3.

Where  $A_0=1, p=1, A_1=\frac{n}{2}+1, B_1=\frac{n}{2}$ .

For p=2

$$A_2^3 + B_2^3 = s_2 (T_3^1 s_2^2 - T_3^2 D_2) = s_2 (s_2^2 - 3D_2) = s_2 \left\{ s_2^2 - 3 \left[ \frac{(s_2 + 1)(s_2 - 1)}{4} - (A_1 - 1)A_1 \right] \right\} = \quad (12)$$

$$= (n^2 + 1) \left\{ (n^2 + 1)^2 - 3 \left[ \frac{(n^2 + 2)n^2}{4} - \left(\frac{n}{2} + 1\right)\frac{n}{2} \right] \right\}$$

For p=3

$$A_3^4 + B_3^4 = (n^3 + 1)^4 - \left\{ \frac{(n^3 + 2)n^3}{4} - \frac{n}{2}(n+1) \left[ \frac{n}{2}(n+1) + 1 \right] \right\} \times$$

$$\times \left\langle 4(n^3 + 1)^2 - 2 \left\{ \frac{(n^3 + 2)n^3}{4} - \frac{n}{2}(n+1) \left[ \frac{n}{2}(n+1) + 1 \right] \right\} \right\rangle \quad (13)$$

### 3. ABIYEV TRIANGLES

As it seen from the formulae, as dimension of space increases, (12) and (13) equalities become more complicated depending on order of geometrical figure. Only p=1, Pythagorean theorem is satisfied. Because, the orders of expressions are the same in both right and left of equality [3].

It should be mentioned that, formulae (6) and (7), which suggested for natural numbers in centres of geometric figures, are true as well as for any numbers.

If p+1=n,  $A_p=a$  and  $B_p=b$  in (6) and (7) equalities and open the brackets, they will be written as follows:

$$a^n + b^n = (a+b) \left( T_n^1 (a+b)^{n-1} - T_n^2 (a+b)^{n-3} (ab) + \dots + T_n^{\frac{n-1}{2}} (a+b)^2 y^{\frac{n-3}{2}} - T_n^{\frac{n+1}{2}} (ab)^{\frac{n-1}{2}} \right) \quad (7')$$

The identity  $a^n \pm b^n = F \left[ (a+b), ab \right]$  was first revealed and a special table was compiled to determine the coefficients in this identity [4]. Let's explain the algorithm first to write the table: insert columns to the right and left of 0 at the top of the table 1 and write numbers beginning from 1 in sequence in column 0. Write odd numbers to the right

side and figure 1 to the left side of this column; write figure 2 before even numbers at the right and figure 0 at the left. In consequence, 2 columns formed at the right and at the left of column 0.

Afterwards, write figure 1 in the 2<sup>nd</sup> and 3<sup>rd</sup> rows of the 2<sup>nd</sup> columns; by summing the figure 1 in the 2<sup>nd</sup> row of the 2<sup>nd</sup> column with the figure 3 in the 3<sup>rd</sup> row of the 1<sup>st</sup> column

write figure 4 in the 4<sup>th</sup> row of the 2<sup>nd</sup> column. So by continuing this method figures 4, 9, 16, 25 etc. squares of sequent numbers are written at the right in the even rows of the 2<sup>nd</sup> column.

Summing the figure 1 in the 3<sup>rd</sup> row of the 2<sup>nd</sup> column with the figure 4 underneath figure 5 is written in the 5<sup>th</sup> row; later summing this last figure 5 with the figure 9 underneath figure 14 is written in the 7<sup>th</sup> row. At the result, the 2<sup>nd</sup> column is formed at the right.

In order to write the 3<sup>rd</sup> column figure 1 is written in the 4<sup>th</sup> and 5<sup>th</sup> rows of the 3<sup>rd</sup> column again; afterwards the above-mentioned method is applied to form the 2<sup>nd</sup> column. So a number of columns and rows of figures can be formed.

**4. IDENTITIES**

Let's accept the following replacements in order to simplify the method of writing polynomial expressions:  $a+b=x$  and is an even power,  $ab=y$ . Let's show the identities of even and odd powers. If nis even power

$$a^n + b^n = E_n^1 x^n - E_n^2 x^{n-2} y + E_n^3 x^{n-4} y^2 - \dots$$

$$\dots - E_n^{\frac{n}{2}} x^2 y^{\frac{n-2}{2}} + E_n^{\frac{n+2}{2}} y^{\frac{n}{2}} \quad (14)$$

here

$$E_n^{k+1} = \frac{n}{k} \binom{n-k-1}{k-1}; \quad k = 0, 1, 2, 3, \dots, \frac{n}{2}; \quad (15)$$

If the method used to form the columns at the right side of the table is applied symmetrically with respect to column 0, columns can be written at the left side too.

The compiled table 1 is composed of right and left triangles. The numbers in the rows of the right and left triangles specify the coefficients of the polynomial expressions to which  $a^n + b^n$  and  $a^n - b^n$  expressions equal respectively. They were put in sequence from right to left in the right triangle and from left to right in the left triangle. One of the interesting properties of this table is that the sum of the numbers at the right and at the left rows forms Lucas and Fibonacci sequences correspondingly.

$$E_n^1 = \frac{n}{0} \binom{n-1}{-1} \equiv 1 \quad \text{is accepted.}$$

If  $n$  is an odd power,

$$a^n + b^n = x \left( T_n^1 x^{n-1} - T_n^2 x^{n-3} y + T_n^3 x^{n-5} y^2 - \dots \right.$$

$$\left. \dots + T_n^{\frac{n-1}{2}} x^2 y^{\frac{n-3}{2}} - T_n^{\frac{n+1}{2}} y^{\frac{n-1}{2}} \right) \quad (16)$$

here

$$T_n^{k+1} = \frac{n}{k} \binom{n-k-1}{k-1}; \quad k = 0, 1, 2, 3, \dots, \frac{n-1}{2} \quad (17)$$

and  $T_n^1 = \frac{n}{0} \binom{n-1}{-1} \equiv 1$  is accepted.

TABLE 2.  
ABIYEV'S TRIANGLES (LEFT AND RIGHT)

	7	6	5	4	3	2	1	0	1	2	3	4	5	6	7	
<b>F</b>								<b>n</b>								<b>L</b>
1							1	1	1							1
1						1	0	2	2	1						3
2						1	1	3	3	1						4
3					1	2	0	4	2	4	1					7
5					1	3	1	5	5	5	1					11
8				1	4	3	0	6	2	9	6	1				18
13				1	5	6	1	7	7	14	7	1				29
21			1	6	10	4	0	8	2	16	20	8	1			47
34			1	7	15	10	1	9	9	30	27	9	1			76

55		1	8	21	20	5	0	10	2	25	50	35	10	1		123
----	--	---	---	----	----	---	---	----	---	----	----	----	----	---	--	-----

The written  $E_n^{k+1}$  and  $T_n^{k+1}$  coefficients can be described in another way too:

$$E_n^{k+1} = \binom{n-k}{k} + \binom{n-1-k}{k-1}; \quad k = 0, 1, 2, \dots, \frac{n}{2} \quad (n \text{-even})$$

$$T_n^{k+1} = \binom{n-k}{k} + \binom{n-1-k}{k-1}; \quad k = 0, 1, 2, \dots, \frac{n-1}{2} \quad (n \text{-odd})$$

Let's write the polynomial expressions for  $a^n - b^n$  expression. If  $n$  is an even power,

$$\frac{a^n - b^n}{a - b} = x \left( M_n^1 x^{n-2} - M_n^2 x^{n-4} y + M_n^3 x^{n-6} y^2 - \dots \dots - M_n^{\frac{n-2}{2}} x^2 y^{\frac{n-4}{2}} + M_n^{\frac{n}{2}} y^{\frac{n-2}{2}} \right) \quad (18)$$

here  $M_n^{k+1} = \binom{n-k-1}{k}; \quad k = 0, 1, 2, 3, \dots, \frac{n}{2} - 1 \quad (19)$

If  $n$  is an odd power,

$$\frac{a^n - b^n}{a - b} = N_n^1 x^{n-1} - N_n^2 x^{n-3} y + N_n^3 x^{n-5} y^2 - \dots \dots + N_n^{\frac{n-1}{2}} x^2 y^{\frac{n-3}{2}} - N_n^{\frac{n+1}{2}} y^{\frac{n-1}{2}} \quad (20)$$

here  $N_n^{k+1} = \binom{n-k-1}{k}; \quad k = 0, 1, 2, 3, \dots, \frac{n-1}{2} \quad (21)$

The written  $M_n^{k+1}$  and  $N_n^{k+1}$  coefficients can be described in another way too:

$$M_n^{k+1} = \binom{n-k}{k} - \binom{n-1-k}{k-1}; \quad k = 0, 1, 2, \dots, \frac{n}{2} - 1 \quad (n \text{-even})$$

$$N_n^{k+1} = \binom{n-k}{k} - \binom{n-1-k}{k-1}; \quad k = 0, 1, 2, \dots, \frac{n-1}{2}; \quad \binom{n-1}{-1} \equiv 0 \quad (n \text{-odd}).$$

Note: As the signs in (14), (16), (18) and (20) polynomial expressions always change in sequence,  $(-1)^n$  symbol hasn't been used in them.

Let's show 2 examples for the identities:

If  $n = 6, \quad a^6 + b^6 = x^6 - 6x^4y + 9x^2y^2 - 2y^3;$

If  $n = 7, \quad \frac{a^7 - b^7}{a - b} = x^6 - 5x^4y + 6x^2y^2 - y^3.$

See the 6<sup>th</sup> row in the right triangle and the 7<sup>th</sup> row in the left triangle of the table 2.

**5. PROPERTIES OF COEFFICIENTS**

It's known that the sum of the numbers in each row of Pascal triangle equals to  $2^{n-1}$ . But in Abiyev's triangles the

sum of the numbers in the rows forms Lucas (at the right) and Fibonacci (at the left) sequences. Though in the literature [5] it's shown that the sum of the numbers in a diagonal line in Pascal triangle forms Fibonacci sequence, what values these numbers have is unknown.

Let's display that in Abiyev's triangles these singularities are not random.

For this purpose the coefficients of (14), (16), (18) and (20) polynomial expressions and Binet formula  $L_n = \alpha^n + \beta^n; F_n = \frac{\alpha^n - \beta^n}{\alpha - \beta}; \alpha = \frac{1+\sqrt{5}}{2}; \beta = \frac{1-\sqrt{5}}{2}$  of Lucas and Fibonacci sequences are used [3]. Here  $x = \alpha + \beta = 1; y = \alpha\beta = -1$ . If these values are put into (14), (16), (18) and (20) polynomial expressions, the following equations can be obtained:

$$a^n + b^n = \alpha^n + \beta^n = E_n^1 + E_n^2 + E_n^3 + \dots + E_n^{\frac{n}{2}} + E_n^{\frac{n+2}{2}} = L_n \quad n \text{-even}; \quad (\text{Lucas term}).$$

$$a^n + b^n = \alpha^n + \beta^n = T_n^1 + T_n^2 + T_n^3 + \dots + T_n^{\frac{n-1}{2}} + T_n^{\frac{n+1}{2}} = L_n; \quad n \text{-odd}; \quad (\text{Lucas term}).$$

$$\frac{a^n - b^n}{a - b} = \frac{\alpha^n - \beta^n}{\alpha - \beta} = M_n^1 + M_n^2 + M_n^3 + \dots + M_n^{\frac{n-2}{2}} + M_n^{\frac{n}{2}} = F_n; \quad n \text{-even}; \quad (\text{Fibonacci term}).$$

$$\frac{a^n - b^n}{a - b} = \frac{\alpha^n - \beta^n}{\alpha - \beta} = N_n^1 + N_n^2 + N_n^3 + \dots + N_n^{\frac{n-1}{2}} + N_n^{\frac{n+1}{2}} = F_n; \quad n \text{-odd}; \quad (\text{Fibonacci term}).$$

So taking into account (15), (17), (19) and (21) polynomial expressions new formulae can be written for Lucas and Fibonacci triangles:

$$L_n = \sum_{k=0}^{\frac{n}{2}} E_n^{k+1} = \sum_{k=0}^{\frac{n}{2}} \frac{n}{k} \binom{n-1-k}{k}; \quad (22a)$$

$$L_n = \sum_{k=0}^{\frac{n-1}{2}} T_n^{k+1} = \sum_{k=0}^{\frac{n-1}{2}} \frac{n}{k} \binom{n-1-k}{k}; \quad (22b)$$

$$F_n = \sum_{k=0}^{\frac{n}{2}-1} M_n^{k+1} = \sum_{k=0}^{\frac{n}{2}-1} \binom{n-1-k}{k}; \quad (23a)$$

$$F_n = \sum_{k=0}^{\frac{n-1}{2}} N_n^{k+1} = \sum_{k=0}^{\frac{n-1}{2}} \binom{n-1-k}{k}; \quad (23b)$$

Let's give examples for  $L_6$  and  $F_7$  formulas:

$$L_6 = \sum_{k=0}^3 E_6^{k+1} = \sum_{k=0}^3 \frac{6}{k} \binom{5-k}{k-1} \Rightarrow E_6^1 + E_6^2 + E_6^3 + E_6^4 = 1 + \frac{6}{1} \binom{4}{0} + \frac{6}{2} \binom{3}{1} + \frac{6}{3} \binom{2}{2} = 1 + 6 + 9 + 2 = 18.$$

$$F_7 = \sum_{k=0}^3 N_7^{k+1} = \sum_{k=0}^3 \binom{6-k}{k} \Rightarrow F_7 = N_7^1 + N_7^2 + N_7^3 + N_7^4 =$$

$$= \binom{6}{0} + \binom{5}{1} + \binom{4}{2} + \binom{3}{3} = 1 + 5 + 6 + 1 = 13.$$

The above-formed formulae enable us to express the following theorem.

**Theorem.** There are Lucas and Fibonacci numbers

$$\text{expressed by } L_n = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n}{k} \binom{n-1-k}{k-1} \text{ and } F_n = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n}{k} \binom{n-1-k}{k}$$

formulas for any positive whole number.

**Proof.** Taking into consideration the peculiarities of the algorithm used for formulating Abiyev's triangles, let's prove the theorem applying a mathematical induction method. According to (15) and (17) formulae let's form right Abiyev's triangle for  $(n-1), n, (n+1), (n+2)$  rows:

In accordance with the singularities of the right and left triangles the following expressions can be written:

$$\text{I} - l_{ij} + l_{i+1,j-1} = l_{i+2,j} ; \quad \text{II} - l_{i-1,j-1} + l_{i,j-1} = l_{i+1,j-1} ; \\ 2 \leq i \leq 2p ; \quad 2 \leq j \leq p+1 ; \quad p \geq 1. \quad (\text{right})$$

$$\text{I} - f_{ij} + f_{i+1,j-1} = f_{i+2,j} ; \quad \text{II} - f_{i-1,j-1} + f_{i,j-1} = f_{i+1,j-1} ; \\ 2 \leq i \leq 2p ; \quad 2 \leq j \leq p+1 ; \quad p \geq 1. \quad (\text{left})$$

here  $l$  and  $f$  are elements of the right and left triangles respectively.

Let's conduct the following algebraic operations using this triangle:

$$\text{I} - l_{n,3} + l_{n+1,2} = l_{n+2,3}$$

$$\frac{n}{2} \binom{\frac{n+2}{2}}{\frac{n-6}{2}} + \frac{n+1}{2} \binom{\frac{n+2}{2}}{\frac{n-4}{2}} = \frac{n+2}{2} \binom{\frac{n+4}{2}}{\frac{n-4}{2}} \Rightarrow$$

$$\Rightarrow \frac{n}{\frac{n-4}{2}} \binom{\frac{n+2}{2}}{4} + \frac{n+1}{\frac{n-2}{2}} \binom{\frac{n+2}{2}}{3} = \frac{n+2}{\frac{n-2}{2}} \binom{\frac{n+4}{2}}{4} \Rightarrow$$

$$\Rightarrow \frac{n}{\frac{n-4}{2}} \frac{\frac{n+2}{2} \frac{n}{2} \frac{n-2}{2} \frac{n-4}{2}}{1.2.3.4} + \frac{n+1}{\frac{n-2}{2}} \frac{\frac{n+2}{2} \frac{n}{2} \frac{n-2}{2}}{1.2.3} =$$

$$= \frac{n+2}{\frac{n-2}{2}} \frac{\frac{n+4}{2} \frac{n+2}{2} \frac{n}{2} \frac{n-2}{2}}{1.2.3.4} \Rightarrow n^2 + 6n + 8 = n^2 + 6n + 8.$$

$$\text{II} - l_{n-1,2} + l_{n,2} = l_{n+1,2}$$

$$\frac{n-1}{\frac{n-4}{2}} \binom{\frac{n}{2}}{\frac{n-6}{2}} + \frac{n}{\frac{n-2}{2}} \binom{\frac{n}{2}}{\frac{n-4}{2}} = \frac{n+1}{\frac{n-2}{2}} \binom{\frac{n+2}{2}}{\frac{n-4}{2}} \Rightarrow$$

$$\Rightarrow \frac{n-1}{\frac{n-4}{2}} \binom{\frac{n}{2}}{3} + \frac{n}{\frac{n-2}{2}} \binom{\frac{n}{2}}{2} = \frac{n+1}{\frac{n-2}{2}} \binom{\frac{n+2}{2}}{3} \Rightarrow$$

$$\Rightarrow \frac{n-1}{\frac{n-4}{2}} \frac{\frac{n}{2} \frac{n-2}{2} \frac{n-4}{2}}{1.2.3} + \frac{n}{\frac{n-2}{2}} \frac{\frac{n}{2} \frac{n-2}{2}}{1.2} =$$

$$= \frac{n+1}{\frac{n-2}{2}} \frac{\frac{n+2}{2} \frac{n}{2} \frac{n-2}{2}}{1.2.3} \Rightarrow n^2 + 3n + 2 = n^2 + 3n + 2.$$

These operations specify that the algorithm of formation in the right triangle is defined for any  $n$ . The same assumption concerns the left triangle too. By this way, the theorem is proved.

TABLE 3. PART OF THE ABIYEV'S RIGHT TRIANGLE

$\frac{col. \rightarrow}{\downarrow row}$	1	2	3	...	$\frac{n-4}{2}$	$\frac{n-2}{2}$	$\frac{n}{2}$	$\frac{n+2}{2}$	$\frac{n+4}{2}$
$n-1:$	$\frac{n-1}{2} \binom{\frac{n-2}{2}}{\frac{n-4}{2}}$	$\frac{n-1}{2} \binom{\frac{n}{2}}{\frac{n-6}{2}}$	$\frac{n-1}{2} \binom{\frac{n+2}{2}}{\frac{n-8}{2}}$	...	$\frac{n-1}{2} \binom{n-4}{1}$	$\frac{n-1}{1} \binom{n-3}{0}$	1		
$n:$	$\frac{n}{2} \binom{\frac{n-2}{2}}{\frac{n-2}{2}}$	$\frac{n}{2} \binom{\frac{n}{2}}{\frac{n-4}{2}}$	$\frac{n}{2} \binom{\frac{n+2}{2}}{\frac{n-6}{2}}$	...	$\frac{n}{3} \binom{n-4}{2}$	$\frac{n}{2} \binom{n-3}{1}$	$\frac{n}{1} \binom{n-2}{0}$	1	
$n+1:$	$\frac{n+1}{2} \binom{\frac{n}{2}}{\frac{n-2}{2}}$	$\frac{n+1}{2} \binom{\frac{n+2}{2}}{\frac{n-4}{2}}$	$\frac{n+1}{2} \binom{\frac{n+4}{2}}{\frac{n-6}{2}}$	...	$\frac{n+1}{3} \binom{n-3}{2}$	$\frac{n+1}{2} \binom{n-2}{1}$	$\frac{n+1}{1} \binom{n-1}{0}$	1	
$n+2:$	$\frac{n+2}{2} \binom{\frac{n}{2}}{\frac{n}{2}}$	$\frac{n+2}{2} \binom{\frac{n+2}{2}}{\frac{n-2}{2}}$	$\frac{n+2}{2} \binom{\frac{n+4}{2}}{\frac{n-4}{2}}$	...	$\frac{n+2}{4} \binom{n-3}{3}$	$\frac{n+2}{3} \binom{n-2}{2}$	$\frac{n+2}{2} \binom{n-1}{1}$	$\frac{n+2}{1} \binom{n}{0}$	1

6. APPLICATIONS

The solution of

$$\begin{cases} X^n \pm Y^n = a \\ X + Y = b \end{cases}$$

equation system is simplified, i.e. the power of the system

is reduced to a half and brought to the solution of  $\frac{n}{2}$  equivalent equation system.

The calculation of  $(X+iY)^n \pm (X-iY)^n = Z^n \pm \bar{Z}^n$  expression mostly used in linear algebra is rather facilitated

For instance, let's show  $C = \left(\frac{1}{2} + i\frac{\sqrt{3}}{2}\right)^7 + \left(\frac{1}{2} - i\frac{\sqrt{3}}{2}\right)^7$ .

Here taking into account

$$x = a + b = 1; \quad y = ab = 1 \quad \text{and}$$

$$a^7 + b^7 = x(x^6 - 7x^4y + 14x^2y^2 - 7y^3),$$

$C = 1 - 7 + 14 - 7 = 1$  is easily calculated.

## 7. CONCLUSION

The content of the article is new. The coefficients of the polynomial expressions can especially be accepted as a spectrum of Lucas and Fibonacci sequences. As these sequences have a significant importance in mathematics, their spectrum can also be applied in different fields. The identities proposed in the article will allow simplifying mathematical calculations in various fields of science and technology.

## REFERENCES

- [1] I.N.Bronstein, K.A.Samandiyev. Handbook on Mathematics, Moscow, Science, Vol.XIII, 1986, p.141 (in Russian);
- [2] A. A. Abiyev, M. Sahin, A. A. Abiyev, A study on Derivation and Explanation of  $A_p^{p+1} + B_p^{p+1} = F\left[\left(A_p + B_p\right), A_p B_p\right]$  Formula For Some Geometric Figures, Journal of Marmara for Pure and Applied Sciences, 18 (2002), 73-80.
- [3] Zhang Zaiming, Problem #502, Pell's Equation-Once Again, Two-year College Math. Inl, 25 (1994), 241-243.
- [4] S.Goktepe, A.A.Abiyev. A New Algorithm to Form Lucas and Fibonacci Sequences, XIX International Symposium on Mathematics, August 22-25, 2006, Kutahya, pp.600- 604 (in Turkish);
- [5] Kenneth H. Rozen, Handbook of Discrete and Combinatorial Mathematics, CRC Press LLC (New York, 2000), pp. 98, 104, 140-143.

**Name:** Asker Ali Abiyev

**Place and date of birth:** Baku, Azerbaijan, 28.06.1934.

**Education:** primary school No 79, and 208 (1944-1954), Baku; Azerbaijan State University (1954-1957); Moscow State University named M. V. Lomonosov, Physic faculty (1957-1961); post graduate student of Institute Atomic Energy named I. V. Kurchatov (1963-1966); Ph.D in physics-mathematics (1970); Doctor in physics-mathematics (1989); Professor (1990).

**Scientific activity:** neutron spectroscopy, radiation physics of semiconductors, magic squares and cubes.

**Career/Employment:**

1.Head of the laboratory radiation physics semiconductors in the Institute of Radiation Problems of Azerbaijan National Academy Sciences 1975-1993; Consultant in the School Boys, Ankara Turkey 1993-2000; Professor of Mathematics in the Gaziantep of University in Turkey, 2000-2007; Head of Experimental Department of the Electron Accelerator, 2009 up today.

**Participated Conferences:** 1. The 2<sup>nd</sup> International Conference on Applied Informatics and Computing Theory (AICT'11), Prague, Czech Republic, September 26-28, 2011.

2.IMS'2008 6th International Symposium on Intelligent and Manufacturing Systems «Feature, Strategies and Innovation», October 14- 17, 2008, Sakarya, Turkey.

3. Fourth International Conference on Soft Computing, Computing with Words and Perceptions in Systems. Analysis, Decision and Control, Turkey, August 27-28, 2007.

4. School of Information Technology and Mathematical Sciences, University Ballarat (Australia). 14 -28 June 2006.

5. Joint International Scientific Conference "New Geometry of Nature", August- September 5,2003, Kazan State University, Kazan, Russia.

6. The Third International Conference On Mathematical And Computational Applications, September 4-6, 2002, Konya,Turkey.

7. 2nd International Conference On Responsive Manufacturing, Gaziantep, Turkey, 26-28, June, 2002.

8. The Second International Symposium on Mathematical and Computational Applications, September 1-3, 1999, Baku, Azerbaijan.



9. Research Conference on Number Theory and Arithmetical Geometry, San Feliu de Guixols, Spain 24-29 October, 1997.  
International Conference on Radiation Physics of Semiconductors and Related Materials, TBİLİSİ, USSR, 1978.

**Publications:** Number of paper in refereed journals: 120  
Number of communications to scientific meetings: 25

**The Book:**

A.K. Abiyev, Sayılı Sihirli Karelerin Doğal Şifresi-The Natural Code of Numbered Magic Squares, Enderun Publications, Ankara, (ISBN 975-95318-3-6), p.77, 1996, (in Turkish and in English ).  
<http://www1.gantep.edu.tr/~bingul/php/magic/>

**Language Skills:** Azeri, Russian, English, Turkish.

**Contact:** askeraliabiye@hotmail.com