

- [5]. Nuriddinova D.Z., Yusupov, F.M. Cleaning water from oil products by carbon mineral sorbent. Volgograd 2018.p.8-11
- [6]. W. Cao, Z. Dang, X.Q. Zhou, X.Y. Yi, P.X. Wu, N.W. Zhu, G.N. Lu, Removal of sulphate from aqueous solution using modified rice straw: preparation, characterization and adsorption performance, Carbohydr. Polym., 2011, vol. 85 p. 571.
- [7].Chinese NEPA, Water and Wastewater Monitoring Methods, (fourth ed.), Chinese Environmental Science Publishing House, Beijing, China, 2012, p. 52.
- [8].UNEP, Water and Wastewater Reuse. An Environmentally Sound Approach for Sustainable Urban Water Management., 2013, USA.
- [9]. Kern-Jespersen J.P, Henze, M, Strube R (1994) Biological Phosphorus Release and Uptake Under Alternating Anaerobic and Anoxic Conditions in a Fixed-film Reactor. Water Research, vol. 28, 1994, p. 1253.
- [10].Asatekin, A.M. Mayes, Oil industry wastewater treatment with fouling resistant membranes containing amphiphilic comb copolymers, Environ. Sci. Technol., Vol. 43, 2009, p. 4487.
- [11].P.D. Fletcher, L.D. Savory, F. Woods, A. Clarke, A.M. Howe, Model study of enhanced oil recovery by flooding with aqueous surfactant solution and comparison with theory, Langmuir, vol. 31, 2015, p. 3076.
- [12]. Praveen K. Bayannavar, Madivalagouda S. Sannaikar, S. Madan Kumar, Sanjeev R. Inamdar, Saba Kauser J. Shaikh, Aravind R. Nesaragi, Ravindra R. Kamble, Synthesis, X-ray characterization, DFT studies and Hirshfeld surface analysis of new organic single crystal: 2-(4-Methoxyphenyl)-4-yl methyl}-2,4-dihydro-3H-1, 2,4-triazol-3-one (MTBT), Journal of Molecular Structure, 2019, 1179: 809-819
- [13]. Thammarat Aree, Suchada Jongrungruangchok, Structure – antioxidant activity relationship of β -cyclodextrin inclusion complexes with olive tyrosol, hydroxytyrosol and oleuropein: Deep insights from X-ray analysis, DFT calculation and DPPH assay, Carbohydrate Polymers, 2018, 199: 661– 669.
- [14].UNEP, Water and Wastewater Reuse. An Environmentally Sound Approach for Sustainable Urban Water Management., 2013, USA.
- [15].Kern-Jespersen J.P, Henze, M, Strube R (1994) Biological Phosphorus Release and Uptake Under Alternating Anaerobic and Anoxic Conditions in a Fixed-film Reactor. Water Research, vol. 28, 1994, p.1253.
- [16].Asatekin, A.M. Mayes, Oil industry wastewater treatment with fouling resistant membranes containing amphiphilic comb copolymers, Environ. Sci. Technol., vol. 43, 2009, p. 4487.
- [17].P.D. Fletcher, L.D. Savory, F. Woods, A. Clarke, A.M. Howe, Model study of enhanced oil recovery by flooding with aqueous surfactant solution and comparison with theory, Langmuir, vol. 31, 2015, p. 3076.
- [18].Praveen K. Bayannavar, Madivalagouda S. Sannaikar, S. Madan Kumar, Sanjeev R. Inamdar, Saba Kauser J. Shaikh, Aravind R. Nesaragi, Ravindra R. Kamble, Synthesis, X-ray characterization, DFT studies and Hirshfeld surface analysis of new organic single crystal: 2-(4-Methoxyphenyl)-4-yl methyl}-2,4-dihydro-3H-1,2,4-triazol-3-one (MTBT), Journal of Molecular Structure, 2019, 1179:809-819
- [19]. Thammarat Aree, Suchada Jongrungruangchok, Structure–antioxidant activity relationship of β -cyclodextrin inclusion complexes with olive tyrosol, hydroxytyrosol and oleuropein: Deep insights from X-ray analysis, DFT calculation and DPPH assay, Carbohydrate Polymers, 2018, 199:661–669.

Creative Commons Attribution License 4.0 (Attribution 4.0 International , CC BY 4.0)

This article is published under the terms of the Creative Commons Attribution License 4.0

https://creativecommons.org/licenses/by/4.0/deed.en_US